

Infection Prevention and Control Practices in Hospitals: A Nursing Perspective

Nitika Thakur

Department of Child Health Nursing Associate Professor, Sharda School of Nursing Science and Research Sharda University, Greater Noida, Uttar Pradesh, India.

Abstract

Background: Infection prevention and control (IPC) is a key component of patient safety and quality care in hospital settings. Healthcare-associated infections (HAIs) remain a major global health challenge, contributing to increased morbidity, mortality, prolonged hospital stays, and higher healthcare costs. Nurses, as frontline healthcare providers, play a central role in implementing IPC practices and reducing infection transmission.

Objectives: This review aims to examine IPC practices in hospitals from a nursing perspective, highlighting key principles, nursing responsibilities, challenges, and strategies for improvement.

Methods: A narrative review of peer-reviewed literature was conducted using electronic databases, including PubMed, Google Scholar, CINAHL, Scopus, and Web of Science. Studies focusing on hospital-based IPC practices, nursing roles, and nursing-led interventions for reducing HAIs were included. Relevant data were organized thematically to summarize IPC principles, hand hygiene practices, use of personal protective equipment, education and training of nurses, and barriers to compliance.

Results: The review reported standard and transmission-based pre-cautions, consistent hand hygiene, proper use of PPE, and continuous nursing education as critical components of effective IPC. Common challenges included workload pressure, limited resources, insufficient training, and variable compliance among nursing staff.

Conclusion: Consistent application of evidence-based IPC measures, supported by ongoing education and institutional commitment, is essential to minimize infection risks. Strengthening nursing involvement in infection prevention strategies can significantly improve patient safety, quality of care, and overall hospital outcomes.

Keywords: Cross infection, hand hygiene, infection control, nursing practice, personal protective equipment

INTRODUCTION

Infection prevention and control (IPC) is a vital component of patient safety and quality care in hospital settings.

Date of Submission: 10-01-2026

Date of Revision: 25-01-2026

Date of Acceptance: 07-02-2026

Access this article online

Website: <http://innovationalpublishers.com/Journal/ijnmi>

ISSN No: 2656-4656

DOI: 10.31690/ijnmi.2026.v01i101.002

Healthcare-associated infections (HAIs) remain a major global public health challenge, contributing to increased morbidity, mortality, prolonged hospital stays, and substantial economic burden on healthcare systems. Hospitals are particularly high-risk environments for infection transmission due to the frequent use of invasive procedures, high patient density, and continuous interaction between healthcare workers and patients.

Effective IPC practices are essential to minimize the risk of infection transmission and ensure safe healthcare delivery. Standard pre-cautions, such as hand hygiene, appropriate use of personal protective equipment (PPE), environmental cleaning, and safe biomedical waste management form the foundation of hospital infection control programs. Adherence

Address for Correspondence:

Nitika Thakur, Department of Child Health Nursing, Associate Professor, Sharda School of Nursing Science and Research Sharda University, Greater Noida, Uttar Pradesh, India.
E-mail: nitithakur.1988@gmail.com

This is an open-access journal, and articles are distributed under the terms of the Creative Commons Attribution Noncommercial Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms

to these practices plays a critical role in protecting patients, healthcare workers, and visitors from preventable infections.

In recent years, the increasing prevalence of antimicrobial resistance and the emergence of new and re-emerging infectious diseases have further highlighted the importance of robust infection prevention strategies. These challenges have emphasized the need for consistent implementation of evidence-based IPC measures and continuous monitoring of compliance in healthcare settings. Strengthening IPC practices is therefore essential for improving patient outcomes, enhancing healthcare quality, and ensuring the overall safety of hospital environments.

Role of nurses in hospital infection control

Nurses play a key role in the successful implementation of IPC practices. As frontline caregivers, nurses are directly involved in patient care activities that carry a high risk of infection transmission. Their responsibilities include maintaining hand hygiene, using PPE correctly, following aseptic techniques, and identifying early signs of infection. Nurses also educate patients and caregivers on infection prevention and contribute to infection surveillance, documentation, and reporting. Their compliance, knowledge, and attitudes significantly influence the effectiveness of IPC programs and patient safety outcomes.^[1]

Aim and objectives of the review

The aim of this review is to examine IPC practices in hospitals from a nursing perspective. The objectives are to describe key IPC principles, explore the role of nurses in infection control, identify common challenges affecting compliance, and highlight strategies to strengthen nursing involvement in reducing HAIS and improving patient outcomes.

METHODOLOGY

Literature search strategy

A comprehensive literature search was conducted to identify relevant studies related to IPC practices in hospital settings from a nursing perspective. Electronic databases, including PubMed, Google Scholar, CINAHL, Scopus, and Web of Science, were searched. Key terms used in various combinations included infection prevention, infection control, HAIS, hospital nursing, nursing practices, and patient safety. Reference lists of selected articles were also reviewed to identify additional relevant studies.^[2]

Inclusion criteria

- Studies focusing on IPC practices in hospital settings
- Articles emphasizing the role of nurses in preventing and controlling HAIS
- Research evaluating nursing-led interventions related to infection control
- Peer-reviewed original research articles, review articles, and clinical guidelines
- Studies published in the English language.

Exclusion criteria

- Studies conducted in non-hospital or community healthcare settings
- Articles not related to nursing practice or nursing involvement in infection control
- Conference abstracts, editorials, opinion pieces, and letters to the editor
- Non-peer-reviewed publications
- Studies published in languages other than English.

Data selection and synthesis approach

Titles and abstracts of retrieved articles were screened for relevance, followed by a full-text review of eligible studies. Relevant data were extracted and organized thematically based on key aspects of IPC practices. A narrative synthesis approach was adopted to summarize findings, identify common themes, and highlight gaps in existing literature.^[3]

OVERVIEW OF HAIS

Definition and types of HAIs

HAIs are infections that develop in patients during hospitalization or while receiving medical care and were not present at the time of admission. These infections typically occur 48 h or more after admission and serve as important indicators of patient safety and quality of healthcare delivery. HAIs are commonly classified based on the site and source of infection.

Major types of HAIs in hospital settings include surgical site infections, catheter-associated urinary tract infections, central line-associated bloodstream infections, ventilator-associated pneumonia, and hospital-acquired gastrointestinal infections, such as *Clostridioides difficile*. These infections are often linked to invasive procedures, prolonged use of medical devices, inadequate hand hygiene, and environmental contamination.

Burden of HAIs in hospital settings

The burden of HAIs in hospitals is considerable, leading to increased patient morbidity and mortality, prolonged hospital stays, and higher healthcare costs. Patients affected by HAIs often require additional treatments and extended antibiotic therapy, which further increases the risk of antimicrobial resistance. HAIs also place a significant strain on healthcare resources and impact the overall quality of care.

Nurses play a vital role in reducing the burden of HAIs through continuous patient monitoring, adherence to infection prevention protocols, and early identification of infection risks. Strengthening nursing-led infection control practices is essential for improving patient safety and reducing infection rates in hospital environments.^[4]

PRINCIPLES OF IPC

Standard pre-cautions

Standard pre-cautions represent the minimum level of infection control practices that should be applied to the care of all

patients, regardless of their suspected or confirmed infection status. These pre-cautions are based on the assumption that blood, body fluids, secretions, excretions, non-intact skin, and mucous membranes may contain potentially infectious microorganisms. Hand hygiene is the cornerstone of standard pre-cautions and includes handwashing with soap and water or the use of alcohol-based hand rubs before and after patient contact.^[5]

The use of PPE, such as gloves, gowns, masks, and eye protection, is another critical component of standard pre-cautions. PPE use depends on the level of anticipated exposure and helps prevent the transmission of pathogens between patients and healthcare workers. Additional elements include safe injection practices, respiratory hygiene and cough etiquette, proper cleaning and disinfection of patient care equipment, and safe handling of linen and biomedical waste. Nurses are responsible for integrating these measures into routine clinical practice to reduce the risk of cross-infection.^[6]

Transmission-based pre-cautions

Transmission-based pre-cautions are additional infection control measures implemented when standard pre-cautions alone are insufficient to prevent the spread of specific infectious agents. These pre-cautions are used for patients with known or suspected infections that are transmitted through contact, droplets, or airborne routes. Contact pre-cautions are employed for infections spread by direct or indirect contact and include the use of gloves, gowns, and dedicated equipment. Droplet pre-cautions are required for pathogens transmitted through respiratory droplets, necessitating the use of surgical masks and spatial separation of patients.^[7]

Airborne pre-cautions are applied to infections transmitted through airborne particles and require specialized measures, such as negative-pressure isolation rooms and the use of respirators. Nurses play a critical role in recognizing the need for transmission-based pre-cautions, educating patients and visitors, and ensuring strict compliance with isolation protocols. Effective implementation of these pre-cautions significantly reduces the risk of infection transmission within hospital settings.

HAND HYGIENE PRACTICES IN NURSING

Hand hygiene is universally recognized as the most effective and fundamental measure for preventing the transmission of infections in hospital settings. From a nursing perspective, hand hygiene plays a critical role in IPC, as nurses have frequent and direct contact with patients, medical devices, and the hospital environment. Proper hand hygiene practices significantly reduce the spread of pathogens, lower the incidence of HAIS, and contribute to improved patient safety and quality of care.^[8]

Importance of hand hygiene

The hands of healthcare workers are a primary route for the transmission of microorganisms between patients and

within the hospital environment. In nursing practice, routine activities, such as patient assessment, wound care, medication administration, and handling of invasive devices create multiple opportunities for microbial transfer. Effective hand hygiene, including handwashing with soap and water or the use of alcohol-based hand rubs, interrupts this transmission pathway and prevents cross-contamination.

International guidelines emphasize the importance of hand hygiene at key moments of patient care, including before and after patient contact, before aseptic procedures, after exposure to body fluids, and after contact with patient surroundings. Consistent adherence to these practices has been shown to reduce rates of HAIS, antimicrobial resistance, and infection-related complications. Nurses, as frontline caregivers, are instrumental in maintaining hand hygiene standards and promoting a culture of safety within healthcare institutions.

Compliance and barriers among nurses

Despite strong evidence supporting hand hygiene, compliance among nurses remains variable across healthcare settings. Factors influencing compliance include workload pressure, staff shortages, time constraints, and high patient acuity. Limited access to hand hygiene supplies, such as alcohol-based hand rubs or handwashing facilities, further contribute to suboptimal adherence. In addition, skin irritation caused by frequent handwashing and lack of ongoing training may discourage consistent practice.

Behavioral and organizational factors also play a role in hand hygiene compliance. Lack of awareness, inadequate monitoring, and insufficient leadership support can negatively impact adherence to protocols.^[9] Addressing these barriers requires a multifaceted approach, including regular education and training programs, availability of resources, institutional policies, and continuous monitoring with feedback. Strengthening nurses' knowledge and commitment to hand hygiene is essential for improving compliance and reducing infection risks in hospital settings.

USE OF PPE

PPE is a critical component of IPC in hospital settings, serving as a barrier between healthcare workers and infectious agents. From a nursing perspective, the appropriate selection and correct use of PPE are essential for preventing the transmission of infections and ensuring the safety of both patients and healthcare personnel.

Types and appropriate use of PPE

- PPE includes gloves, gowns, masks, respirators, eye protection, and face shields
- Each type of PPE is designed to protect against specific routes of infection transmission, such as contact, droplet, or airborne exposure
- The selection of appropriate PPE depends on the nature of patient care, the level of exposure risk, and the suspected or confirmed mode of transmission

- Proper donning (putting on) and doffing (removing) techniques are essential to prevent self-contamination and cross-infection
- Nurses must follow established PPE guidelines during routine patient care activities
- PPE use is particularly important during high-risk procedures, including handling body fluids, performing invasive procedures, and caring for patients with communicable diseases
- Adherence to PPE protocols helps protect both healthcare workers and patients and supports effective IPC.

Nursing responsibilities

- Ensure correct selection and appropriate use of PPE based on patient condition and risk of exposure
- Follow institutional infection control guidelines and standard operating procedures related to PPE use
- Practice proper donning and doffing techniques to prevent self-contamination
- Educate patients and visitors on the importance and correct use of PPE when required
- Maintain compliance with PPE use during routine care and high-risk procedures
- Report shortages or defects in PPE to hospital administration to ensure safety.

Challenges in PPE use

- Shortages or limited availability of PPE, especially during outbreaks or high patient load
- Physical discomfort and fatigue associated with prolonged use of PPE
- Increased workload and time constraints affecting consistent PPE compliance
- Inadequate training or lack of regular updates on correct PPE practices
- Communication difficulties with patients while wearing masks or face shields.

EDUCATION AND TRAINING OF NURSES

Education and training are fundamental pillars for the effective implementation of IPC practices in hospital settings. Nurses, as the largest group of healthcare professionals and primary providers of direct patient care, play a critical role in preventing HAIs. Their knowledge, clinical skills, and attitudes toward infection control directly influence patient safety, quality of care, and overall health outcomes. Adequate education ensures that nurses are well equipped to apply evidence-based IPC measures consistently and appropriately across diverse clinical situations. Formal nursing education provides the foundation for infection control knowledge, including microbiology, principles of asepsis, and standard pre-cautions. However, initial education alone is insufficient to meet the dynamic challenges of modern healthcare. Continuous training and professional development are essential to ensure that nurses remain competent, confident, and responsive to evolving

infection risks, emerging pathogens, and updated national and international guidelines.^[10]

Infection control training programs

Infection control training programs are essential for equipping nurses with the theoretical knowledge and practical skills required to prevent, identify, and manage HAIs. These programs commonly address key components, such as standard and transmission-based pre-cautions, hand hygiene practices, appropriate use of PPE, safe injection practices, environmental cleaning, sterilization and disinfection procedures, and proper segregation and disposal of biomedical waste. Structured infection control training is particularly important for newly recruited nurses, as early reinforcement of correct practices helps establish safe clinical behaviors and professional accountability. Orientation programs that include IPC protocols reduce practice variability and promote uniform adherence to institutional policies. For experienced nurses, periodic refresher training ensures retention of knowledge and correction of unsafe or outdated practices. Regular in-service education sessions, workshops, and simulation-based learning approaches enhance practical understanding and encourage active participation. Simulation training, in particular, allows nurses to practice infection control measures in realistic clinical scenarios, improving confidence and decision-making skills. Infection control training also plays a crucial role during outbreaks, pandemics, and public health emergencies by rapidly updating nurses on emerging infectious diseases, revised guidelines, and emergency preparedness protocols.^[11]

Effective training programs contribute to improved compliance with infection prevention measures, reduction in HAIs, and development of a strong culture of safety within healthcare institutions. Administrative support, availability of resources, and multidisciplinary collaboration further strengthen the impact of these training initiatives.^[12]

Continuing education and competency development

Continuing education is essential for maintaining and enhancing nurses' competence in IPC throughout their professional careers. Rapid advancements in medical technology, evolving infection control guidelines, increasing antimicrobial resistance, and the emergence of novel pathogens necessitate lifelong learning among nursing professionals. Continuing professional development programs, online learning modules, seminars, and certification courses provide opportunities for nurses to stay updated with present best practices. Competency development involves regular assessment of both knowledge and practical skills related to infection control. Key competencies include hand hygiene compliance, correct donning and doffing of PPE, maintenance of aseptic techniques, safe management of invasive devices, and adherence to isolation protocols. Periodic audits, skill demonstrations, and performance evaluations help identify gaps and areas requiring improvement. Constructive feedback, mentorship, and supportive supervision play a vital role in

reinforcing correct practices and promoting accountability. Continuous monitoring and competency-based training foster professional growth and encourage nurses to take ownership of infection prevention responsibilities. Investing in continuing education and competency development not only strengthens nursing practice but also leads to sustained reductions in HAIS, improved patient outcomes, and enhanced quality of hospital care.^[13,14]

CONCLUSION

IPC practices remain a cornerstone of patient safety and quality healthcare delivery in hospital settings. Nurses, as frontline healthcare providers with continuous patient contact, play a pivotal role in minimizing the risk of HAIS through diligent hand hygiene, correct use of PPE, strict adherence to standard and transmission-based pre-cautions, and active participation in infection surveillance, reporting, and patient education. Their role extends beyond task-based compliance to influencing safety culture and modeling best practices within multidisciplinary healthcare teams. Despite the availability of well-established national and international IPC guidelines, consistent implementation continues to face significant challenges. Factors, such as high patient workload, staff shortages, limited access to infection control resources, inadequate infrastructure, and gaps in ongoing training negatively impact adherence to recommended practices. These challenges underscore the need for sustained institutional commitment rather than reliance on individual responsibility alone. Addressing these barriers requires a comprehensive and systematic approach that includes regular education and training, competency-based assessments, supportive supervision, and strong organizational leadership. Continuous professional development programs and routine monitoring can enhance knowledge retention, skill proficiency, and accountability among nursing staff. Furthermore, ensuring the availability of essential resources, including PPE, hand hygiene facilities, and infection surveillance systems, is critical for enabling effective practice. Strengthening nursing-led infection control initiatives, supported by hospital management and policy frameworks, can significantly contribute to the reduction of HAIS. Such efforts not only improve patient outcomes and reduce morbidity and mortality but also enhance overall healthcare quality, reduce healthcare costs, and promote safer hospital environments. Sustained investment in nursing education, workforce support, and infection control infrastructure is therefore essential for achieving long-term improvements in IPC outcomes.

CONFLICT OF INTEREST

NA.

SOURCE OF FUNDING

NA.

ACKNOWLEDGMENT

NA.

REFERENCES

1. Kubde D, Badge AK, Ugemuge S, Shahu S. Importance of hospital infection control. *Cureus* 2023;15:e50931.
2. Eriksen MB, Frandsen TF. The impact of patient, intervention, comparison, outcome (PICO) as a search strategy tool on literature search quality: A systematic review. *J Med Libr Assoc* 2018;106:420-31.
3. Raab GM, Nowok B, Dibben C. Practical data synthesis for large samples. *J Priv Confid* 2016;7:67-97.
4. Gidey K, Gidey MT, Hailu BY, Gebreamlak ZB, Niriayo YL. Clinical and economic burden of healthcare-associated infections: A prospective cohort study. *Plos One* 2023;18:e0282141.
5. Curran ET. Standard precautions: What is meant and what is not. *J Hosp Infect* 2015;90:10-1.
6. Reda AA, Fisseha S, Mengistie B, Vandeweerd JM. Standard precautions: Occupational exposure and behavior of health care workers in Ethiopia. *PLoS One* 2010;5:e14420.
7. Link T. Guideline implementation: Transmission-based precautions. *AORN J* 2019;110:637-49.
8. Labrague LJ, McEnroe-Petitte DM, Van De Mortel T, Nasirudeen AM. A systematic review on hand hygiene knowledge and compliance in student nurses. *Int Nurs Rev* 2018;65:336-48.
9. Alhumaid S, Al Mutair A, Al Alawi Z, Alsuliman M, Ahmed GY, Rabaan AA, *et al.* Knowledge of infection prevention and control among healthcare workers and factors influencing compliance: A systematic review. *Antimicrob Resist Infect Control* 2021;10:86.
10. Zhao Y, Xu Y, Ma D, Fang S, Zhi S, He M, *et al.* The impact of education/training on nurses caring for patients with stroke: A scoping review. *BMC Nurs* 2024;23:90.
11. Tsioutis C, Birgand G, Bathoorn E, Deptula A, Ten Horn L, Castro-Sánchez E, *et al.* Education and training programmes for infection prevention and control professionals: Mapping the current opportunities and local needs in European countries. *Antimicrob Resist Infect Control* 2020;9:183.
12. Tartari E, Fankhauser C, Masson-Roy S, Márquez-Villarreal H, Fernández Moreno I, Rodríguez Navas ML, *et al.* Train-the-trainers in hand hygiene: A standardized approach to guide education in infection prevention and control. *Antimicrob Resist Infect Control* 2019;8:206.
13. Cox DR, Grus CL. From continuing education to continuing competence. *Prof Psychol Res Pract* 2019;50:113-9.
14. Salum NC, Prado ML. Continuing education in the development of competences in nurses. *Texto Contexto Enferm* 2014;23:301-8.

How to cite this article: Thakur N. Infection Prevention and Control Practices in Hospitals: A Nursing Perspective. *Int J Nurs Med Invest.* 2026;11(1):6-10.